

## A Case of Methotrexate-Associated Lymphoproliferative Disorder After Resection of an Accidentally Detected Enlarged Thyroid Tumor

Masatsugu Yano<sup>1\*</sup>, Daisuke Sumitani<sup>1</sup>, Masayuki Shishida<sup>1</sup>, Takafumi Oshiro<sup>1</sup>, Makoto Ochi<sup>1</sup>, Yuzo Okamoto<sup>1</sup>, Hirofumi Nakayama<sup>2</sup> and Hideki Ohdan<sup>3</sup>

<sup>1</sup>Department of Surgery, JR Hiroshima Hospital, Hiroshima, Japan

<sup>2</sup>Department of Pathology and Laboratory Medicine, JR Hiroshima Hospital, Hiroshima, Japan

<sup>3</sup>Department of Gastroenterological and Transplant Surgery, Applied Life Sciences, Institute of Biomedical and Health Sciences, Hiroshima University, Hiroshima, Japan

### \*Corresponding author:

Masatsugu Yano,  
Department of Surgery, JR Hiroshima Hospital 3-1-36,  
Futabanosato, Higashi-ku, Hiroshima 732-0057,  
Japan, Tel: +81-082-262-1170;  
Fax: +81-082-262-1499;  
E-mail: yano4masa@marble.ocn.ne.jp

Received: 09 Nov 2021

Accepted: 16 Nov 2021

Published: 22 Nov 2021

J Short Name: ACMCR

### Copyright:

©2021 Masatsugu Yano. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

### Citation:

Masatsugu Yano, A Case of Methotrexate-Associated Lymphoproliferative Disorder After Resection of an Accidentally Detected Enlarged Thyroid Tumor. *Ann Clin Med Case Rep.* 2021; V7(16): 1-5

### Keywords:

Methotrexate-associated lymphoproliferative disorders (MTX-LPD); Thyroid; Rheumatoid arthritis

## 1. Abstract

**1.1. Introduction:** We herein present a case of methotrexate-associated lymphoproliferative disorder (MTX-LPD) with a diagnosis of an 8-mm thyroid tumor of class III cytology that grew three times larger during 2 years 8 months before resection.

**1.2. Case Presentation:** The patient was a 71-year-old woman who was receiving methotrexate treatment for rheumatoid arthritis. A thyroid tumor was identified on computed tomography scans taken to closely examine the abnormal shadows observed on chest radiographs taken during a routine health checkup. She was then referred to our department. Ultrasound of the thyroid confirmed an 8-mm tumor in the right lobe of the thyroid, and the patient was considered for a class III cytology diagnosis. Owing to the enlargement of the tumor at 2 years 8 months after the first examination, the patient underwent surgery. Histopathological examination revealed that the tumor was a diffuse large B-cell lymphoma; given the patient's use of methotrexate, it was conceived as MTX-LPD. Discontinuing MTX treatment and observing the patient's clinical course is considered the first course of action to be taken for MTX-LPD.

**1.3. Conclusion:** This is a rare case of MTX-LPD that was confirmed while monitoring a microthyroid tumor discovered accidentally and operated on before the manifestation of MTX-LPD symptoms. Radical excisional biopsy could be performed even before the patient shows symptoms such as neck swelling.

## 2. Introduction


Methotrexate (MTX) is a drug that is often administered to patients with rheumatoid arthritis. At times, patients develop lymphoproliferative disorder during the course of treatment, and the condition is referred to as MTX-associated lymphoproliferative disorder (MTX-LPD). Here we describe a case of MTX-LPD involving an 8-mm thyroid tumor of class III cytological diagnosis, and this tumor grew three times larger during 2 years 8 months before resection.

## 3. Case Report


A 71-year-old woman was receiving methotrexate treatment for rheumatoid arthritis. Chest radiographs were taken during a routine health checkup. Later, she was referred to our department and computed tomography (CT) scans were performed to closely examine the abnormal shadows observed on the chest radiographs,

which revealed a thyroid tumor. There were no palpable masses in the thyroid or cervical lymph nodes. Ultrasound examination revealed an 8-mm hypoechoic tumor in the right lobe of the thyroid (Figure 1). Fine needle aspiration cytology revealed a Bethesda category III finding. Therefore we decided to follow up the tumor. Two years eight months after the first examination, the tumor grew threefold, to a size of 24 mm over (Figure 2). We elected to perform surgery as a suspicion of malignancy. Then she underwent surgery (right-sided hemithyroidectomy) and was discharged on postoperative day 7 on account of a favorable clinical course. Histopathological examinations were performed on a tumor specimen with


dimensions of 25X20X20mm. Hematoxylin-eosin (HE) staining showed proliferation of lymphoma cells with multiple nucleoli in a large nucleus. There were multiple mitotic presentations, and the cells lacked cytoplasm (Figure. 3). Immunostaining showed CD20 positivity, CD3 negativity, and an MIB-1 index of approximately 50% (Figure. 4). The histopathological presentations of the tumor specimen were similar to those of diffuse large B-cell lymphoma. We surmised that she had developed MTX-LPD due to her use of MTX for treating chronic rheumatoid arthritis. After a 2-year follow-up, local recurrence was not detected.




**Figure 1:** Thyroid ultrasonography image showing an 8-mm hypoechoic tumor in the right lobe of the thyroid.




**Figure 2:** Image showing the threefold growth of the tumor, from 8 mm to 24 mm, over 2 years 8 months since the initial examination.




**Figure 3:** Image of the resected tumor specimen with dimensions of 25X20X20mm (A). Hematoxylin-eosin staining image showing the proliferation of lymphoma cells with multiple nucleoli in a large nucleus. There are multiple mitotic presentations and the cells lack cytoplasm. (B).



**CD3**



**CD20**



**MIB-1**

**Figure 4:** Immunostaining image showing CD20 positivity, CD3 negativity, and the MIB-1 index was approximately 50%.

#### 4. Discussion and Conclusion

MTX has been administered as an anticancer drug for various types of carcinomas such as leukemia and breast cancer. It is also administered as an immunosuppressant for autoimmune disease such as rheumatoid arthritis and dermatomyositis. MTX-LPD was first reported in 1991 by Ellman et al. based on their experience with a patient with rheumatoid arthritis who was taking oral

MTX and who developed lymphoma and their report discussed the possibility of immunosuppression by MTX contributing to the development of lymphoma [1]. In 1993, Kamel et al. reported two cases of lymphoma that developed during oral MTX treatment [2], and this report was supported by multiple similar reports in the following years [3-6]. Currently, the World Health Organization classifies MTX-LPD as an “immunodeficiency-associated lymphoproliferative disease” and considers it as a pathology caused by immunodeficiency [7]. Hoshida et al. reported that the median age at onset of MTX-LPD diagnosis was 67 years, the male-to-female ratio of patients with this condition was approximately 1:2 the mean duration from RA onset to MTX-LPD onset was 11 years, and the mean duration of MTX treatment was 5 years [3]. Furthermore Yoshihara et al. reported that the mean age at MTX-LPD onset was 71 years, the mean duration of MTX treatment was 9 years. However, the duration of treatment was not necessarily long, with an even distribution across patients from 0.3 years to 17 years [4]. Kameda et al. analyzed several clinical parameters and reported that the dose of MTX used is a risk factor [8]. Yoshihara et al. strongly suspected a link between high-dose MTX and the number of MTX-LPD cases [4]. However, no clear risk factor for MTX-LPD is yet to be reported. It is said that the duration of MTX treatments, total dose of MTX used, or severity of RA cannot be considered risk factors [4]. Compared to other lymphomas, lymphomas due to MTX-LPD tend to manifest more frequently as extranodal lesions and are common in the gastrointestinal tract, skin, lungs, and soft tissues. In the head and neck regions, such lymphomas are seen in the cervical lymph nodes and tonsils; as extranodal lesions, they have been seen in the salivary glands, thyroid glands, and tooth buds [3, 6, 9-12]. The histopathological presentations of MTX-LPD are diverse, with the most common diffuse large B-cell lymphoma accounting for 35%-60% of cases, followed by Hodgkin lymphoma and follicular lymphoma, which account for 12%-25% and 5%-10%, respectively [3-6, 10, 13]. There are reports suggesting the involvement of the Epstein-Barr (EB) virus as a cause of the disease, with 32%-71% of MTX-LPD cases tested positive for EB virus [2, 14, 15]. According to the disease type, it has been reported that approximately 80% of Hodgkin lymphoma cases and approximately 20% of diffuse large B-cell lymphoma cases are positive for EB virus [5]. Rheumatoid arthritis patients are reported to have a 2- to 20-fold increased risk of developing lymphoma even without MTX treatment [16]. Patients treated with MTX are reported to be at an even higher risk of developing lymphoma, as the treatment changes the balance between EB virus load and the host [17]. Discontinuing MTX treatment and observing the patient’s clinical course is considered the first course of action to be taken for MTX-LPD, after confirming its histological diagnosis [4, 6, 18-20]. If there is tumor regression, the patient’s condition needs to be monitored carefully. However, discontinuation of MTX treatment alone after MTX-LPD onset lead to com-

plete remission in 11 out of 48 cases as reported Hoshida et al. [3], in 3 out of 8 cases as reported by Mariette et al. [5], and in 3 out of 13 cases as reported by Kojima et al [10]. If discontinuing MTX treatment is not enough on its own to see clinical remission, patients should be provided treatment according the type of malignant lymphomas [4]. The five-year survival rate of MTX-LPD is reportedly more than 70%, but patient’s age of more than 70 years old and pathological subtype of DLBCL are known to be indicators of a poor prognosis [21-22]. Our patients had an 8-mm thyroid tumor with a class III cytological diagnosis that was detected accidentally, and we decided to observe the patient’s clinical course. The tumor grew in size by three times during the course (2 years 8 months) after the initial examination. Histopathological examination of the tumor specimen after hemithyroidectomy suggested a diagnosis of MTX-LPD. This is a rare case for which radical excisional biopsy could be performed even before the patient shows symptoms such as neck swelling. After a 2-year follow-up, local recurrence was not detected. It would be necessary to keep the patient under careful observation.

If a thyroid tumor is identified in a patient who is treated with oral MTX, then the possibility of MTX-LPD should be taken into consideration. It is also important to monitor the patient for the presence of nodules in the thyroid gland, even if it is so small.

## References

1. Ellman MH, Hurwitz H, Thomas C, Kozloff M. Lymphoma developing in a patient with rheumatoid arthritis taking low dose weekly methotrexate. *J Rheumatol.* 1991; 18: 1741-1743.
2. Kamel OW, van de Rijn M, Weiss LM, Zoppo GJD, Hench PK, Rossins BA, et al. Brief report: reversible lymphomas associated with Epstein-Barr virus occurring during methotrexate therapy for rheumatoid arthritis and dermatomyositis. *N Engl J Med.* 1993; 328 (18): 1317-1321.
3. Hoshida Y, Xu JX, Fujita S, Nakamichi I, Ikeda JI, Tomita Y, et al. Lymphoproliferative disorders in rheumatoid arthritis: clinicopathological analysis of 76 cases in relation to methotrexate medication. *J Rheumatol.* 2007; 34 (2): 322-331.
4. Yoshihara R, Uefuji A, Murata M, Nakagawa N, Tanaka Y, Shiozawa K. Clinical manifestations of twenty-seven cases of methotrexate-associated lymphoproliferative disorders. *Clin Rheumatol Rel Res* 2017; 29: 164-172.
5. Mariette X, Cazals-hatem D, Warszawski J, Liote F, Balandraud N, Sibilia J, et al. Lymphomas in rheumatoid arthritis patients treated with methotrexate. A 3-yeara prospective study in France. *Blood.* 2002; 99 (11): 3909-3915.
6. Salloum E, Cooper DL, Howe G, Lacy J, Tallini G, Crouch J, et al. Spontaneous regression of lymphoproliferative disorders in patients treated with methotrexate for rheumatoid arthritis and other rheumatic disease. *J Clin Oncol.* 1996; 14 (6): 1943-1949.
7. Gaulard P, Swerdlow SH, Harris NL, Sundstrom C, Jaffe ES.

- Other iatrogenic immunodeficiency associated lymphoproliferative disorders. WHO classification of tumors of haemtopoietic and lymphoid tissues. IARC Press Lyon. 2017: 462-464.
8. Kameda T, Dobashi H, Miyatake N, Inoo M, Onishi I, Kurata N, et al. Association of higher methotrexate dose with lymphoproliferative disease onset in rheumatoid arthritis patients. *Arthritis Care Res.* 2014; 66 (9): 1302-1309.
  9. Ishida Y, Asahi A, Wada T, Kanai N, Kobayashi Y, Moriai S, et al. Three cases of methotrexate-associated lymphoproliferative disorder. *Nippon Jibiinkoka Gakkai Kaiho.* 2008; 111: 594-598.
  10. Kojima M, Itoh H, Hirabayashi K, Igarashi S, Tamaki Y, Murayama K, et al. Methotrexate-associated lymphoproliferative disorders. A clinicopathological study of 13 Japanese cases. *Pathl Res Pract* 2006; 202 (9): 679-685.
  11. Feng WH, Cohen JI, Fisher S, Li L, Sneller M, Goldbach-Mansky R. Reactivation of latent Epstein-Barr virus by methotrexate: a potential contributor to methotrexate-associated lymphoma. *J Natl Cancer Inst.* 2004; 96 (22): 1691-1702.
  12. Lim IGS, Bertouch JV. Remission of lymphoma after drug withdrawal in rheumatoid arthritis. *Med J Aust.* 2002; 177 (9): 500-501.
  13. Baird RD, van Zyl-Smit RN, Dilke T, Scott SE, Rassam SMB. Spontaneous remission of low-grade B-cell non-Hodgkin's lymphoma following withdrawal of methotrexate in a patient with rheumatoid arthritis: case report and review of the literature. *Br J Haematol.* 2002; 118 (2): 567-568.
  14. Kawano N, Ono N, Yoshida S, Kuriyama T, Yamashita K, Beppu K, et al. Successful treatment of immunodeficiency-associated EBV-negative lymphoproliferative disorders in rheumatoid arthritis by methotrexate withdrawal and prevention of its relapse by rituximab administration. *J Clin Exp Hematopathol.* 2012; 52 (3): 193-198.
  15. Callan M. Epstein-Barr virus, arthritis, and the development of lymphoma in arthritis patients. *Cur Opin Rheumatol.* 2004; 16 (4): 399-405.
  16. Tokuhira M, Watanabe R, Nemoto T, Sagawa M, Tomikawa T, Tamaru J. Clinicopathological analyses in patients with other iatrogenic immunodeficiency –associated lymphoproliferative diseases and rheumatoid arthritis. *Leuk Lymphoma.* 2012; 53 (4): 616-623.
  17. Ichikawa A, Arakawa F, Kiyasu J, Sato K, Miyoshi H, Niino D, et al. Methotrexate/iatrogenic lymphoproliferative disorders in rheumatoid arthritis: histology, Epstein-Barr virus, and clonarity are important predictors of disease progression and regression. *Eur J Haematol.* 2013; 91 (1): 20-28.
  18. Ishida M, Hodohara K, Yoshii M, Okuno H, Horinouchi A, Nakanishi R, et al. Methotrexate-related Epstein-Barr virus-associated lymphoproliferative disorder occurring in the gingiva of a patients with rheumatoid arthritis. *Int J Clin Exp Pathol.* 2013; 6 (10): 2237-2241.
  19. Thomas E, Brewster DH, Black RJ, Macfarlane GJ. Risk of malignancy among patients with rheumatic conditions. *Int J Cancer.* 2000; 88 (3): 497-502.
  20. Koens L, Senff N, Vermeer M, Willemze R, Jansen P. Methotrexate-associated B-cell lymphoproliferative disorders presenting in the skin: A clinicopathologic and immunophenotypical study of 10 cases. *Am J Surg Pathol.* 2014; 38 (7): 999-1006.
  21. Ohkura Y, Shindoh J, Haruta S, Kajii D, Ota Y, Fujii T, et al. Primary adrenal lymphoma possibly associated with Epstein-Barr virus reactivation due to immunosuppression under methotrexate therapy. *Medicine.* 2015; 94 (31): e1270.
  22. Loo EY, Medeiros LJ, Aladily TN, Hoehn D, Kanagal-Shamanna R, Young KH, et al. Classical Hodgkin lymphoma arising in the setting of iatrogenic immunodeficiency: a clinicopathologic study of 10 cases. *Am J Surg Pathol.* 2013; 37 (8): 1290-1297.