

Disseminated Infection Characterization On FDG-PET/CT: Clinico-Laboratory radiological Correlation Unraveling The Complexity

Tanvi Sarwal¹, Sameer Taywade^{2*}, Althaf K Majeed³, Rajesh Kumar⁴ and Bharat Kumar⁵

¹Junior Resident, Department of Nuclear Medicine, AIIMS Jodhpur, Rajasthan, India.

²Associate Professor, Department of Nuclear Medicine, AIIMS Jodhpur, Rajasthan, India.

³Senior Resident, Department of Nuclear Medicine, AIIMS Jodhpur, Rajasthan, India.

⁴Professor and Head, Department of Nuclear Medicine, AIIMS Jodhpur, Rajasthan, India.

⁵Associate Professor, Department of Medicine, AIIMS Jodhpur, Rajasthan, India.

Corresponding author:

Sameer Taywade,

Associate Professor, Department of Nuclear Medicine, AIIMS Jodhpur, Rajasthan, India.

Phone number: +918130535917

Received Date: 12 Sep 2024

Accepted Date: 01 Oct 2024

Published Date: 07 Oct 2024

Citation:

Sameer Taywade. Disseminated Infection Characterization On FDG-PET/CT: Clinico-Laboratory radiological Correlation Unraveling The Complexity. Annals of Clinical and Medical Case Reports 2024.

1. Abstract

¹⁸F FDG PET/CT is a hybrid imaging technique traditionally used for various oncological indications. Recent advances have broadened its use to various non-oncological indications, such as infections and inflammatory diseases. While FDG uptake indicates increased metabolic activity, however the uptake of FDG is not specific, and it is not possible to distinguish separate infections on the basis of FDG uptake alone. We present a case, wherein through critical analysis of PET/CT images, and correlation with patient's clinical and laboratory data, we were able to identify three distinct infections on FDG PET/CT.

Keywords:

¹⁸F FDG PET/CT, infections, micro-organisms.

2. Introduction

Infections pose a serious threat globally due to the increasing problem of antibiotic resistance [1]. Early diagnosis or exclusion of infection is of utmost importance, and it is a major factor that determines the patient's prognosis [2]. In most cases, the diagnosis of infections is straightforward, using clinical and laboratory data, supplemented with structural imaging techniques such as ultrasonography, computed tomography and magnetic resonance imaging. However, they may not be able to localize the infectious foci in all cases [3]. Thus, functional and metabolic imaging techniques are often needed to complement the role of anatomical imaging methods in many clinical situations [3]. ¹⁸F fluorodeoxyglucose positron emission tomography/ computed tomography (¹⁸F FDG PET/CT) is an advanced hybrid imaging technique that combines the anatomic information derived from CT with the functional information derived from PET. Initially limited to oncological indications, FDG PET/CT has expanded its scope to include non-oncological applications, such as imaging infections and inflammation as well. It allows for whole-body examination in a single session, to identify infection foci in patients [4, 5, 6, 7]. White blood cells have a high glucose metabolism compared to other cells. Additionally, inflammatory mediators cause an upregulation of glucose transporters, leading to increased FDG uptake. Therefore, sites of infection often light up on FDG PET/CT [8]. However FDG is considered relatively nonspecific due its limitations in differentiating infection from inflammation, as well as distinguishing different types of infections. We present an interesting case of a patient with a complex medical history, wherein FDG PET/CT helped to identify the sites of infection which further correlated with three distinct types of organisms.

3. Materials and methods

A 53 year old gentleman presented to the emergency with multiple painful skin eruptions over the body. On examination, the patient was vitally stable and febrile to touch. There were multiple tender non-itchy pustular skin lesions over the scalp, face, neck and chest with a ~2 cm palpable left inguinal lymph node. Co-morbidities included type 2 diabetes mellitus and hypertension. His past history was significant for genitourinary tuberculosis, for which the patient received a course of anti-tubercular therapy, followed by augmentation ileo-cystoplasty for thimble bladder two months back. There was no history of immunosuppressant intake. Routine laboratory workup demonstrated raised inflammatory markers, with elevated total leukocyte count and neutrophils. Blood cultures were sent and empirical antibiotics were started. 2-D echocardiography and ¹⁸F FDG PET/CT were done to identify the primary source of infection.

Annals of Clinical and Medical Case Reports

Echocardiography was negative for vegetations on heart valves. Further to rule out any possibility of myocardial involvement and to look for focus of active infection, patient was referred to 18F FDG PET/CT. The patient was prepared according to infective endocarditis protocol and was instructed to follow a high fat and low carbohydrate diet for 2 days before the PET/CT scan [9]. 18F FDG PET/CT images were acquired from vertex to mid thigh after 55 minutes of injection of 6.5 millicuries of 18F FDG (Figure 1, A). Dedicated cardiac spot images were also acquired. 18F FDG PET/CT revealed multiple metabolically active variable sized skin and subcutaneous nodular cystic lesions involving face, scalp, head & neck, anterior chest & abdominal wall, back, pelvic region and scrotum (SUVmax - 6.6 in the lesion on anterior abdominal wall, Figure 1, B). In addition, there was a thick walled cavitary lesion involving the right lung upper lobe, with peripheral FDG uptake (SUVmax - 5.0). This lesion demonstrated internal necrosis and air fluid levels within (Figure 1, C). Further, there were multiple parenchymal foci of increased FDG uptake involving bilateral renal parenchyma, persistent till delayed images (left kidney, SUVmax - 9.5, delayed SUVmax - 12.9, Figure 1, D). Moreover, irregular focal bladder wall thickening was noted involving the neo-bladder with few hypermetabolic areas (SUVmax - 18.0, delayed SUVmax - 25.3, Figure 1, E). No abnormal FDG uptake was noted in the myocardium.




Figure 1: 18F FDG PET/CT images of the patient. A: Maximum intensity projection image, depicting multiple FDG avid oval lesions, predominantly involving the thorax (solid arrow), as well as multiple foci of increased radiotracer uptake involving bilateral kidneys (dotted arrow). B-D: Fused 18F FDG PET/CT images of the patient. B: FDG avid skin and subcutaneous lesions involving thorax and bilateral axilla. C: Peripherally FDG avid cavitary lesion involving the right lung upper lobe. D: Multiple foci of FDG uptake involving both kidneys. Ancillary evidence of right simple renal cyst. E: Focal hypermetabolism involving the thickened urinary bladder wall.

We tried to correlate the 18F FDG PET/CT findings with the clinical details, laboratory parameters and follow-up imaging of the patient. The biopsy from left inguinal lymph node was suggestive of budding yeasts with pseudohyphae, and culture of pustular skin lesions came out to be positive for *Candida albicans*. This confirmed the fungal origin

of cutaneous lesions seen on 18F FDG-PET/CT. Thus, a diagnosis of disseminated Candidemia was made. Cutaneous tuberculosis was ruled out, since skin lesions were negative for acid fast bacilli. Secondly, the thick walled cavitary lesion in right lung upper lobe was initially suspected to be an aspergilloma based on the CT features [10], however a negative galactomannan assay in bronchoalveolar lavage fluid ruled out this possibility. Further, the presence of a solitary well-defined cavitary lesion with normal remaining bilateral lung parenchyma made pulmonary tuberculous involvement less likely [11]. The patient was started on intravenous caspofungin as per the culture and sensitivity reports, followed by maintenance therapy with oral posaconazole. The skin lesions gradually improved. Further, a follow-up thorax CT scan performed after three months of anti-fungal therapy showed a 50% reduction in the size of the lung lesion. This finding corroborated the possible fungal nature of this lung lesion. Thirdly, *Escherichia coli* (*E. coli*) was identified on urine culture, and ultrasonography of the kidneys demonstrated bilateral pyelonephritis. This helped in correlating metabolically active renal parenchymal lesions to pyelonephritis secondary to *E. coli* infection. Lastly, the focal hypermetabolic bladder wall thickening was correlated with the reactivation of genitourinary tuberculosis, in view of recent urine Cartridge-Based Nucleic Acid Amplification Test (CBNAAT) positivity.

4. Results

In this case, there were distinct infective foci demonstrated on 18F FDG PET/CT involving the skin, lung, kidneys and urinary bladder by three different causative organisms. These included *Candida albicans* in skin lesions, bilateral renal parenchymal lesions attributed to *E. coli*, and reactivation of genitourinary tuberculosis in the form of bladder involvement by *Mycobacterium tuberculosis*. Moreover, the patient presented with superficial *Candida* skin lesions, and FDG PET/CT helped in detecting systemic involvement, specifically in the form of the cavitary lung lesion. This is clinically relevant, as systemic candidiasis requires more aggressive and prolonged antifungal therapy compared to localized skin infections [12].

5. Discussion and Conclusion

Infectious and inflammatory diseases are usually characterized by a more diffuse and less pathognomonic pattern of FDG uptake than FDG uptake for oncological indications. In addition, patients who are referred for FDG PET/CT with suspected infection or inflammation are rarely treatment naive, and may have already received varying dose of antibiotics [13]. Consequently, FDG PET/CT is generally considered as nonspecific modality to distinguish infections [3, 14, 15]. However, critical analysis of the images along with clinical, laboratory and radiological correlation can widen the scope of FDG PET/CT beyond its established applications and help in correlating with different infectious processes occurring simultaneously. There is paucity of data on the role of FDG PET/CT in fungal infections. Our findings align with previous research [16, 17], which has demonstrated a potential role of FDG PET/CT in diagnosing invasive

Annals of Clinical and Medical Case Reports

fungal infections. Moreover, although FDG PET/CT is often considered as less useful for diagnosing tuberculosis of the urinary tract due to the excretory pathway of FDG [18], a careful examination of images, along with correlation with patient's history, can help overcome this limitation.

6. Acknowledgement

The authors are grateful to the patient for agreeing to provide the valuable case for publication.

7. Consent from Patient

The patient has agreed to use his images for publication and provided written informed consent for the same.

References

- Ventola, C. L. The antibiotic resistance crisis: part 1: causes and threats. *Pharmacy and therapeutics*, (2015); 40(4), 277.
- Setyawati, R., Astuti, A., Utami, T. P., Adiwijaya, S., & Hasyim, D. M. The importance of early detection in disease management. *Journal of World Future Medicine, Health and Nursing*, (2024); 2(1), 51-63.
- Gotthardt, M., Bleeker-Rovers, C. P., Boerman, O. C., & Oyen, W. J. Imaging of inflammation by PET, conventional scintigraphy, and other imaging techniques. *Journal of Nuclear Medicine*, (2010); 51(12), 1937-1949.
- Pijl, J. P., Londema, M., Kwee, T. C., Nijsten, M. W., Slart, R. H., Dierckx, R. A., ... & Pillay, J. FDG-PET/CT in intensive care patients with bloodstream infection. *Critical Care*, 25, 1-12. Vaidyanathan S, Patel CN, Scarsbrook AF, Chowdhury FU. FDG PET/CT in infection and inflammation--current and emerging clinical applications. *Clin Radiol*. (2021); 2015 Jul; 70(7):787-800. doi: 10.1016/j.crad.2015.03.010. Epub 2015 Apr 25. PMID: 25917543.
- Vaidyanathan, S., Patel, C. N., Scarsbrook, A. F., & Chowdhury, F. U. FDG PET/CT in infection and inflammation—current and emerging clinical applications. *Clinical radiology*, (2015); 70(7), 787-800.
- Casali, M., Lauri, C., Altini, C., Bertagna, F., Cassarino, G., Cistaro, A., ... & Signore, A. State of the art of 18F-FDG PET/CT application in inflammation and infection: a guide for image acquisition and interpretation. *Clinical and Translational Imaging*, (2021); 9(4), 299-339.
- Jamar, F., Buscombe, J., Chiti, A., Christian, P. E., Delbeke, D., Donohoe, K. J., ... & Signore, A. EANM/SNMMI guideline for 18F-FDG use in inflammation and infection. *Journal of Nuclear Medicine*, (2013); 54(4), 647-658.
- Pijl, J. P., Kwee, T. C., Slart, R. H., & Glaudemans, A. W. PET/CT imaging for personalized management of infectious diseases. *Journal of personalized medicine*, (2021); 11(2), 133.
- Ten Hove, D., Slart, R. H. J. A., Sinha, B., Glaudemans, A. W. J. M., & Budde, R. P. J. 18F-FDG PET/CT in infective endocarditis: indications and approaches for standardization. *Current cardiology reports*, (2021); 23(9), 130.
- Greene, R. The radiological spectrum of pulmonary aspergillosis. *Medical mycology*, (2005); 43(Supplement_1), S147-S154.
- Jeong, Y. J., & Lee, K. S. Pulmonary tuberculosis: up-to-date imaging and management. *American Journal of Roentgenology*, (2008); 191(3), 834-844.
- Pappas, P. G., Rex, J. H., Sobel, J. D., Filler, S. G., Dismukes, W. E., Walsh, T. J., & Edwards, J. E. Guidelines for treatment of candidiasis. *Clinical infectious diseases*, (2004); 38(2), 161-189.
- Pijl, J. P., Nienhuis, P. H., Kwee, T. C., Glaudemans, A. W., Slart, R. H., & Gormsen, L. C. Limitations and pitfalls of FDG-PET/CT in infection and inflammation. In *Seminars in nuclear medicine* (2021, November); (Vol. 51, No. 6, pp. 633-645). WB Saunders.
- Chang, J. M., Lee, H. J., Goo, J. M., Lee, H. Y., Lee, J. J., Chung, J. K., & Im, J. G. False positive and false negative FDG-PET scans in various thoracic diseases. *Korean Journal of Radiology*, (2006); 7(1), 57-69.
- Singh, S. B., Bhandari, S., Siwakoti, S., Bhatta, R., Raynor, W. Y., Werner, T. J., ... & Revheim, M. E. Is imaging bacteria with PET a realistic option or an illusion?. *Diagnostics*, (2023); 13(7), 1231.
- Vadi, S., Kumar, R., Parihar, A., Singh, H., & Mittal, B. in evaluation of disease extent and response evaluation post therapy in invasive fungal diseases: A tertiary care experience (2020) 18F-FDG PET/CT.
- Sharma, P., Mukherjee, A., Karunanithi, S., Bal, C., & Kumar, R. Potential role of 18F-FDG PET/CT in patients with fungal infections. *American Journal of Roentgenology*, (2014); 203(1), 180-189.
- Harkirat, S., Anana, S. S., Indrajit, L. K., & Dash, A. K. Pictorial essay: PET/CT in tuberculosis. *The Indian journal of radiology & imaging*, (2008); 18(2), 141.